


# ABL2 Antibody (R32426)


| Catalog No. | Formulation   | Size   |
|-------------|---|--------|
| R32426      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

# **Bulk quote request**


| Availability       | 1-3 business days  |
|--------------------|--|
| Species Reactivity | Human, Mouse, Rat  |
| Format             | Antigen affinity purified  |
| Clonality          | Polyclonal (rabbit origin)   |
| Isotype            | Rabbit IgG   |
| Purity             | Antigen affinity   |
| Buffer             | Lyophilized from 1X PBS with 2% Trehalose                          |
| UniProt            | P42684   |
| Localization       | Cytoplasmic, membranous, nuclear                                   |
| Applications       | Western Blot : 0.1-0.5ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml |
| Limitations        | This ABL2 antibody is available for research use only.             |



IHC testing of FFPE human gallbladder carcinoma tissue with ABL2 antibody. HEIR: Boil the paraffin sections in pH8 EDTA buffer for 20 minutes and allow to cool prior to testing.



IHC testing of FFPE mouse brain tissue with ABL2 antibody. HEIR: Boil the paraffin sections in pH8 EDTA buffer for 20 minutes and allow to cool prior to testing.



Western blot testing of 1) rat brain, 2) rat PC-12, 3) mouse brain, 4) mouse NIH 3T3 and 5) human K562 cell lysate with ABL2 antibody. Expected molecular weight ~128 kDa, but routinely observed at 128-140 kDa.

## **Description**

Tyrosine-protein kinase ABL2, also known as Abelson-related gene (Arg), is an enzyme that in humans is encoded by the ABL2 gene. This gene encodes a member of the Abelson family of nonreceptor tyrosine protein kinases. The protein is highly similar to the c-abl oncogene 1 protein, including the tyrosine kinase, SH2 and SH3 domains, and it plays a role in cytoskeletal rearrangements through its C-terminal F-actin- and microtubule-binding sequences. This gene is expressed in both normal and tumor cells, and is involved in translocation with the ets variant 6 gene in leukemia. Multiple alternatively spliced transcript variants encoding different protein isoforms have been found for this gene.

### **Application Notes**

Optimal dilution of the ABL2 antibody should be determined by the researcher.

#### **Immunogen**

Amino acids T1065-R1182 of the human protein were used as the immunogen for the ABL2 antibody.

#### **Storage**

After reconstitution, the ABL2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.