


ACTN4 Antibody / Alpha Actinin 4 (R32459)


Catalog No.	Formulation	Size
R32459	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	O43707
Localization	Cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This ACTN4 antibody is available for research use only.




Western blot testing of 1) rat skeletal muscle, 2) rat liver, 3) rat lung, 4) rat RH35, 5) mouse skeletal muscle, 6) mouse liver, 7) mouse lung and 8) mouse NIH 3T3 cell lysate with ACTN4 antibody at 0.5ug/ml. Predicted molecular weight ~105 kDa.




Western blot testing of human 1) MCF7, 2) HeLa, 3) HepG2, 4) HT1080, 5) placenta, 6) A549, 7) U-251 MG and 8) RT-4 cell lysate with ACTN4 antibody at 0.5ug/ml. Predicted molecular weight ~105 kDa.




IHC staining of FFPE human colonic adenocarcinoma tissue with ACTN4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human prostate cancer tissue with ACTN4 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human HeLa cells with ACTN4 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Flow cytometry testing of human U-87 MG cells with ACTN4 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= ACTN4 antibody.

Description

Alpha-actinin-4 is a protein that in humans is encoded by the ACTN4 gene. Alpha actinins belong to the spectrin gene superfamily which represents a diverse group of cytoskeletal proteins, including the alpha and beta spectrins and dystrophins. Alpha actinin is an actin-binding protein with multiple roles in different cell types. In nonmuscle cells, the cytoskeletal isoform is found along microfilament bundles and adherens-type junctions, where it is involved in binding actin to the membrane. In contrast, skeletal, cardiac, and smooth muscle isoforms are localized to the Z-disc and analogous dense bodies, where they help anchor the myofibrillar actin filaments. This ACTN4 gene encodes a nonmuscle, alpha actinin isoform which is concentrated in the cytoplasm, and thought to be involved in metastatic processes. Mutations in this gene have been associated with focal and segmental glomerulosclerosis.

Application Notes

Optimal dilution of the ACTN4 antibody should be determined by the researcher.

Immunogen

Amino acids E561-V661 from the human protein were used as the immunogen for the ACTN4 antibody.

Storage

Prior to reconstitution, store at 4oC. After reconstitution, the ACTN4 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.