

## Adiponectin Antibody (R31122)


Catalog No.	Formulation	Size
R31122	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide and 0.025% thimerosal
<b>UniProt</b>	Q15848
<b>Applications</b>	Western Blot : 0.5-1ug/ml IHC (FFPE) : 0.5-1ug/ml IHC (Frozen) : 0.5-1ug/ml ELISA : 0.1-0.5ug/ml (human protein tested)
<b>Limitations</b>	This Adiponectin antibody is available for research use only.




Western blot testing of Adiponectin antibody and rat heart tissue lysate.  
Expected/observed size ~30KD




Western blot testing of Adiponectin antibody and recombinant protein; Lane 1: 10ng; 2: 5ng; 3: 2.5ng




IHC-P: Adiponectin antibody testing of human placenta tissue



IHC-P: Adiponectin antibody testing of mouse kidney tissue



IHC-P: Adiponectin antibody testing of rat kidney tissue



IHC-F testing of Adiponectin antibody and rat liver tissue

## Description

Adiponectin, also called ADIPOQ (Adipocyte-, C1q-, and Collagen Domain-Containing), APM1, ADPN and ACDC, is a protein which in humans is encoded by the ADIPOQ gene. Using FISH, Das et al.(2001) mapped the mouse Acrp30 gene to chromosome 16 in a region showing homology of synteny with human 3q27. By RNase protection and Western blot analysis, Schaffler et al.(1999) showed that ADIPOQ is expressed by differentiated adipocytes as a 30-kD protein that is also detectable in serum. By sequence comparisons, they found links between ADIPOQ and TNF family ligands as well as to cytokines expressed by T cells. It is a protein hormone that modulates a number of metabolic processes, including glucose regulation and fatty acid oxidation. It is exclusively secreted from adipose tissue (and also from the placenta in pregnancy) into the bloodstream and is very abundant in plasma relative to many hormones.

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the Adiponectin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

An amino acid sequence from the C-terminus of human ADIPOQ (LYADNDNDSTFTGFLLYHDTN) was used as the immunogen for this Adiponectin antibody.

## Storage

After reconstitution, the Adiponectin antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.