


ADPRC1 Antibody / ADP-ribosyl cyclase 1 / CD38 [clone CD38/9333] (V5827)


Catalog No.	Formulation	Size
V5827-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5827-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5827SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	CD38/9333
Purity	Protein G affinity
UniProt	P28907
Localization	Membrane
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 2-4ug/ml
Limitations	This ADPRC1 antibody is available for research use only.




Western blot testing of human Raji tissue lysate with ADPRC1 antibody (clone CD38/9333). Expected molecular weight: 34-46 kDa depending on glycosylation level.



IHC staining of FFPE human tonsil tissue with ADPRC1 antibody (clone CD38/9333).
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human tonsil tissue with ADPRC1 antibody (clone CD38/9333).
Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free ADPRC1 antibody (clone CD38/9333) as confirmation of integrity and purity.

Description

CD38/ADPRC1 is a type II transmembrane glycoprotein that is present on early B- and T-cell lineages and activated B- and T-cells but is absent from most mature resting peripheral lymphocytes. CD38 is also found on thymocytes, pre-B cells, germinal center B-cells, mitogen-activated T-cells, monocytes and Ig-secreting plasma cells. CD38 is expressed on CD34+ cells. The CD34+CD38- population of hematopoietic stem cells define the most pluripotent cells (e.g. blast colony forming cells).

Application Notes

Optimal dilution of the ADPRC1 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-200 from human CD38 protein was used as the immunogen for the ADPRC1 antibody.

Storage

Aliquot the ADPRC1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

