



Alpha-1-Antichymotrypsin Antibody / AACT / SERPINA3 [clone SERPINA3/4189] (V9484)

Catalog No.	Formulation	Size
V9484-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9484-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9484SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	SERPINA3/4189
Purity	Protein A/G affinity
UniProt	P01011
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Alpha-1-Antichymotrypsin antibody is available for research use only.





IHC staining of FFPE human liver tissue with Alpha-1-Antichymotrypsin antibody (clone SERPINA3/4189). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Alpha-1-Antichymotrypsin antibody (clone SERPINA3/4189) as confirmation of integrity and purity.

Description

It recognizes a protein of 65-76kDa, which is identified antichymotrypsin (AACT). AACT is a plasma protease inhibitor synthesized in the liver as a single glycopeptide chain. In human, the normal serum level of AACT is about one-tenth that of their concentrations in plasma increase in response to trauma, surgery and infection. Elevated levels of AACT are widely, but not universally, reported in the cerebrospinal fluid and plasma of AD patients. Prostate-specific antigen (PSA) and its SDS-stable complex with AACT are in widespread use as markers for the diagnosis of prostate cancer. AACT deficiency may also be a possible cause of chronic liver disease. AACT antibody reacts with histiocytes and histiocytic neoplasms. It is widely used to identify histiocytes and tumors derived from them. Acinar tumors of the pancreas and salivary gland may also exhibit AACT positivity.

Application Notes

Optimal dilution of the Alpha-1-Antichymotrypsin antibody should be determined by the researcher.

Immunogen

A portion of amino acids A portion of amino acids 49-187 was used as the immunogen for the Alpha-1-Antichymotrypsin antibody.

Storage

Aliquot the Alpha-1-Antichymotrypsin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.