


Alpha-Synuclein Antibody / SNCA [clone AGA-19] (RQ8886)

Catalog No.	Formulation	Size
RQ8886	Antibody in PBS with 0.02% sodium azide, 50% glycerol and 0.4-0.5mg/ml BSA	100 ul


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	AGA-19
Purity	Affinity chromatography
UniProt	P37840
Localization	Nuclear, cytoplasmic, cell membrane
Applications	Western Blot : 1:500-1:2000 Immunohistochemistry (FFPE) : 1:50-1:200 Immunofluorescence : 1:50-1:200
Limitations	This Alpha-Synuclein antibody is available for research use only.




IHC staining of FFPE human colon cancer tissue with Alpha-Synuclein antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human glioma tissue with Alpha-Synuclein antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse brain tissue with Alpha-Synuclein antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE mouse brain tissue with Alpha-Synuclein antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE rat brain tissue with Alpha-Synuclein antibody, HRP-secondary and DAB substrate. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




Immunofluorescent staining of FFPE human HeLa cells with Alpha-Synuclein antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Immunofluorescent staining of FFPE human HeLa cells with Alpha-Synuclein antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Immunofluorescent staining of FFPE human HeLa cells with Alpha-Synuclein antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of human 1) HEL, 2) RT4 and 3) HeLa cell lysate with Alpha-Synuclein antibody. Predicted molecular weight ~15 kDa.