


Anti-p53 Antibody [clone SPM590] (V9090)


Catalog No.	Formulation	Size
V9090-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V9090-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V9090SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V9090IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b kappa
Clone Name	SPM590
Purity	Protein G affinity chromatography
UniProt	P04637
Localization	Predominantly nuclear
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This anti-p53 antibody is available for research use only.



Western blot testing of human HeLa cell lysate with anti-p53 antibody (clone SPM590).
Expected molecular weight ~53 kDa.



IHC: Formalin-fixed, paraffin-embedded human colon carcinoma stained with anti-p53 antibody (clone SPM590).

Description

Recognizes a 53kDa protein, which is identified as p53 suppressor gene product. It reacts with the mutant as well as the wild form of p53. p53 is a tumor suppressor gene expressed in a wide variety of tissue types and is involved in regulating cell growth, replication, and apoptosis. It binds to MDM2, SV40 T antigen and human papilloma virus E6 protein. Positive nuclear staining with p53 antibody has been reported to be a negative prognostic factor in breast carcinoma, lung carcinoma, colorectal, and urothelial carcinoma. Anti-p53 positivity has also been used to differentiate uterine serous carcinoma from endometrioid carcinoma as well as to detect intratubular germ cell neoplasia. Mutations involving p53 are found in a wide variety of malignant tumors, including breast, ovarian, bladder, colon, lung, and melanoma.

Application Notes

The optimal dilution of the anti-p53 antibody for each application should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human wild type p53 protein was used as the immunogen for this anti-p53 antibody. Its epitope maps within the N-terminus (approx. amino acids 37-45).

Storage

Store the anti-p53 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).