


APOBEC3G Antibody [clone 6C2] (RQ5219)


Catalog No.	Formulation	Size
RQ5219	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ul

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1
Clone Name	6C2
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	Q9HC16
Applications	Western Blot : 0.1-0.5ug/ml Immunohistochemistry(FFPE) : 0.5-1ug/ml Flow Cytometry : 1-3ug/1x10 ⁶ cells Immunofluorescence : 2-4ug/ml
Limitations	This APOBEC3G antibody is available for research use only.




Immunofluorescent staining of FFPE human MCF7 cells with APOBEC3G antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.




IHC staining of FFPE human tonsil with APOBEC3G antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human testis cancer with APOBEC3G antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human intestinal cancer with APOBEC3G antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) Raji and 2) K562 cell lysate with APOBEC3G antibody. Predicted molecular weight ~46 kDa.



Flow cytometry testing of human ThP-1 cells with APOBEC3G antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= APOBEC3G antibody.

Description

APOBEC3G (apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3G) is a human enzyme encoded by the APOBEC3G gene. This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. It is thought that the proteins may be RNA editing enzymes and have roles in growth or cell cycle control. The protein encoded by this gene has been found to be a specific inhibitor of human immunodeficiency virus-1 (HIV-1) infectivity.

Application Notes

Optimal dilution of the APOBEC3G antibody should be determined by the researcher.

Immunogen

A recombinant human protein (amino acids E191-N384) was used as the immunogen for the APOBEC3G antibody.

Storage

After reconstitution, the APOBEC3G antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.