


# **Apolipoprotein A I Antibody (R32177)**


Catalog No.	Formulation	Size
R32177	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

# **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Mouse
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
UniProt	Q00623
Localization	Cytoplasmic, membranous
Applications	Western Blot : 0.1-0.5ug/ml IHC (FFPE) : 0.5-1ug/ml
Limitations	This Apolipoprotein A I antibody is available for research use only.



Western blot testing of mouse liver lysate with Apolipoprotein A I antibody. Predicted molecular weight ~30 kDa.



IHC testing of FFPE mouse liver with Apolipoprotein A I antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.



IHC testing of FFPE mouse kidney with Apolipoprotein A I antibody. HIER: Boil the paraffin sections in pH 6, 10mM citrate buffer for 20 minutes and allow to cool prior to staining.

## **Description**

Apolipoprotein A-1, also known as APOA1, is a human protein with a specific role in lipid metabolism. It binds to lipopolysaccharide or endotoxin, and has a major role in the anti-endotoxin function of HDL. The ApoA-I protein promotes cholesterol efflux from tissues to the liver for excretion. It is a cofactor for lecithin cholesterolacyltransferase (LCAT) which is responsible for the formation of most plasma cholesteryl esters. ApoA-I is also isolated as a prostacyclin (PGI2) stabilizing factor, and thus may have an anticlotting effect. Defects in the gene encoding it are associated with HDL deficiencies, including Tangier disease, and with systemic non-neuropathic amyloidosis. Additionally, ApoA-I overexpression promotes macrophage-specific reverse cholesterol transport.

#### **Application Notes**

Optimal dilution of the Apolipoprotein A I antibody should be determined by the researcher.

#### **Immunogen**

Amino acids 25-264 of mouse Apolipoprotein A I were used as the immunogen for the Apolipoprotein A I antibody.

### **Storage**

After reconstitution, the Apolipoprotein A I antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.