


AR Antibody (F42684)

Catalog No.	Formulation	Size
F42684-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F42684-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Predicted Reactivity	Primate
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity
UniProt	P10275
Applications	Western Blot : 1:1000
Limitations	This AR antibody is available for research use only.



AR antibody western blot analysis in (1) NCI-H292, and (1) NCI-H460 lysate. Predicted molecular weight ~99kDa.

Description

The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract causes spinal bulbar muscular atrophy (Kennedy disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Two

alternatively spliced variants encoding distinct isoforms have been described. [provided by RefSeq].

Application Notes

Titration of the AR antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 282-309 from the human protein was used as the immunogen for this AR antibody.

Storage

Aliquot the AR antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.