


ARHGEF7 Antibody / Beta-Pix / Rho guanine nucleotide exchange factor 7 (FY12862)


Catalog No.	Formulation	Size
FY12862	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml	100 ug

[Bulk quote request](#)


Availability	1-2 days
Species Reactivity	Human, Mouse, Rat
Format	Lyophilized
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Immunogen affinity purified
Buffer	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
UniProt	Q14155
Localization	Cytoplasm
Applications	Western Blot : 0.25-0.5ug/ml Immunohistochemistry : 2-5ug/ml Immunocytochemistry : 5ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells ELISA : 0.1-0.5ug/ml
Limitations	This ARHGEF7 antibody is available for research use only.




Immunofluorescent staining of FFPE human colon cancer tissue with ARHGEF7 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH8 EDTA buffer for 20 min.




Immunofluorescent staining of FFPE human prostate cancer tissue with ARHGEF7 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH8 EDTA buffer for 20 min.




Immunohistochemical staining of ARHGEF7 using anti-ARHGEF7 antibody. ARHGEF7 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARHGEF7 antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Immunohistochemical staining of ARHGEF7 using anti-ARHGEF7 antibody. ARHGEF7 was detected in a paraffin-embedded section of human lung adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARHGEF7 antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Immunohistochemical staining of ARHGEF7 using anti-ARHGEF7 antibody. ARHGEF7 was detected in a paraffin-embedded section of human lung adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARHGEF7 antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Immunohistochemical staining of ARHGEF7 using anti-ARHGEF7 antibody. ARHGEF7 was detected in a paraffin-embedded section of human prostate adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARHGEF7 antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.




Immunohistochemical staining of ARHGEF7 using anti-ARHGEF7 antibody. ARHGEF7 was detected in a paraffin-embedded section of human prostate adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARHGEF7 antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.



Immunohistochemical staining of ARHGEF7 using anti-ARHGEF7 antibody. ARHGEF7 was detected in a paraffin-embedded section of human rectum adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARHGEF7 antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.



Immunohistochemical staining of ARHGEF7 using anti-ARHGEF7 antibody. ARHGEF7 was detected in a paraffin-embedded section of human rectum adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARHGEF7 antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.



Immunohistochemical staining of ARHGEF7 using anti-ARHGEF7 antibody. ARHGEF7 was detected in a paraffin-embedded section of human testicular seminoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ARHGEF7 antibody overnight at 4oC. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37oC. The tissue section was developed using an HRP secondary and DAB substrate.

Description

ARHGEF7 antibody detects Rho guanine nucleotide exchange factor 7, a cytoskeletal signaling protein that regulates Rho GTPase activation, cell adhesion, and migration. Encoded by the ARHGEF7 gene on chromosome 13q34, this protein-also known as Beta-PIX (PAK-interacting exchange factor beta)-functions as a guanine nucleotide exchange factor (GEF) for the small GTPases Rac1 and Cdc42. Through these interactions, ARHGEF7 controls actin cytoskeleton remodeling, focal adhesion turnover, and lamellipodia formation essential for cell motility and morphology.

Structurally, ARHGEF7 contains several modular domains, including an SH3 domain that interacts with PAK kinases, a DH (Dbl homology) domain responsible for nucleotide exchange activity, and a PH (pleckstrin homology) domain that binds phosphoinositides for membrane targeting. These domains enable ARHGEF7 to act as a scaffold connecting small GTPases, kinases, and adhesion proteins such as GIT1 and paxillin. It localizes predominantly at focal adhesions and synaptic sites, integrating extracellular signals with intracellular cytoskeletal responses.

The ARHGEF7 antibody is widely used in cell signaling, cancer biology, and neurobiology research to study cytoskeletal organization, cell motility, and synaptic regulation. Western blot analysis detects a 76 kilodalton band corresponding to ARHGEF7, while immunofluorescence reveals punctate staining along the cell cortex, focal adhesions, and neuronal synapses. This antibody supports exploration of actin dynamics and Rho GTPase signaling in both normal and pathological contexts.

Functionally, ARHGEF7 interacts with p21-activated kinases (PAKs) to form complexes that control Rac1/Cdc42 activation and downstream signaling to the MAPK and JNK pathways. In neurons, it contributes to dendritic spine formation and synaptic plasticity by linking postsynaptic density proteins to actin assembly machinery. Dysregulation of ARHGEF7 has been associated with cancer metastasis, psychiatric disorders, and developmental defects due to altered Rho signaling. The ARHGEF7 antibody provides a dependable tool for dissecting these pathways and quantifying GEF expression in various models. NSJ Bioreagents validates this antibody for its applications, ensuring robust performance

for cytoskeletal and signaling studies.

Application Notes

Optimal dilution of the ARHGEF7 antibody should be determined by the researcher.

Immunogen

E.coli-derived human ARHGEF7 recombinant protein (Position: S25-N637) was used as the immunogen for the ARHGEF7 antibody.

Storage

After reconstitution, the ARHGEF7 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.