


ATRX Antibody [clone ATRX/7188R] (V4021)

Catalog No.	Formulation	Size
V4021-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4021-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4021SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	ATRX/7188R
Purity	Protein A/G affinity
UniProt	P46100
Localization	Nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This ATRX antibody is available for research use only.



IHC staining of FFPE human colon tissue with ATRX antibody (clone ATRX/7188R).
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

ATRX is a member of the Snf2 family of helicase/ATPases, which contribute to the remodeling of the nucleosome structure in an ATP-dependent manner, and facilitate the initiation of transcription and replication. Structurally, ATRX contains a PHD zinc finger motif. ATRX is regulated throughout the cell cycle where it is differentially distributed within the nucleus. During interphase, ATRX predominately associates with the nuclear matrix, while during mitosis, ATRX localizes

with condensed chromatin. At the onset of M phase, phosphorylation rapidly induces this redistribution of ATRX to the short arms of human acrocentric chromosomes, where it then specifically complexes with heterochromatin protein 1
thalassemia or ATRX syndrome

Application Notes

Optimal dilution of the ATRX antibody should be determined by the researcher.

Immunogen

A portion of amino acids 2200-2450 from the human protein was used as the immunogen for the ATRX antibody.

Storage

Aliquot the ATRX antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.