


BATF2 Antibody [clone PCRP-BATF2-2B9] (V4174)


Catalog No.	Formulation	Size
V4174-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4174-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4174SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	PCRP-BATF2-2B9
Purity	Protein A/G affinity
UniProt	Q8N1L9
Localization	Nucleus
Applications	ELISA (Order BSA-free Format For Coating) : Flow Cytometry : 1-2ug/million cells Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This BATF2 antibody is available for research use only.




Flow cytometry testing of PFA-fixed human HeLa cells with BATF2 antibody (clone PCRP-BATF2-2B9) followed by goat anti-mouse IgG-CF488 (blue); isotype control (red).



SDS-PAGE analysis of purified, BSA-free BATF2 antibody (clone PCRP-BATF2-2B9) as confirmation of integrity and purity.



IHC staining of FFPE human prostate carcinoma tissue with BATF2 antibody (clone PCRP-BATF2-2B9). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Analysis of a HuProt(TM) microarray containing more than 19,000 full-length human proteins using BATF2 antibody (clone PCRP-BATF2-2B9). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a mAb to its intended target. A mAb is considered to specific to its intended target, if the mAb has an S-score of at least 2.5. For example, if a mAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that mAb to protein X is equal to 29.

Description

Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II cis-regulatory region sequence-specific DNA binding activity. Predicted to be involved in defense response to protozoan; myeloid dendritic cell differentiation; and regulation of transcription by RNA polymerase II. Predicted to be part of chromatin. Predicted to be active in nucleus. This antibody recognizes a transcription factor involved in differentiation of CD8+ thymic dendritic cells. BATF2 has been implicated in breast cancer, malignant glioma and metastasis melanoma progression.

Application Notes

Optimal dilution of the BATF2 antibody should be determined by the researcher.

Immunogen

A recombinant partial protein (within amino acids 5-140) from the human protein was used as the immunogen for the BATF2 antibody.

Storage

Aliquot the BATF2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

