


Beta Tubulin Antibody [clone 2E11.] (RQ5619)


Catalog No.	Formulation	Size
RQ5619	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a
Clone Name	2E11.
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P07437
Applications	Western Blot : 0.5-1ug/ml Flow Cytometry : 1-3ug/10 ⁶ cells Immunocytochemistry : 2-4ug/ml
Limitations	This Beta Tubulin antibody is available for research use only.




IF/ICC staining of FFPE human U-2 OS cells with Beta Tubulin antibody (green) at 2ug/ml and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.




IF/ICC staining of FFPE human U-2 OS cells with Beta Tubulin antibody (green) at 2ug/ml and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.




Western blot testing of 1) human HEK293, 2) monkey COS-7, 3) human PC-3 and 4) human HeLa cell lysate with Beta Tubulin antibody. Predicted molecular weight: 50-55 kDa.



Western blot testing of rat 1) brain, 2) liver, 3) PC-12 and mouse 4) brain, 5) NIH3T3 and 6) RAW264.7 cell lysate with Beta Tubulin antibody. Predicted molecular weight: 50-55 kDa.



Flow cytometry testing of human U937 cells with Beta Tubulin antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Beta Tubulin antibody.



Flow cytometry testing of human HEPA1-6 cells with Beta Tubulin antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Beta Tubulin antibody.

Description

Tubulin beta chain is a protein that in humans is encoded by the TUBB gene. This gene encodes a beta tubulin protein. This protein forms a dimer with alpha tubulin and acts as a structural component of microtubules. Mutations in this gene cause cortical dysplasia, complex, with other brain malformations 6. Alternative splicing results in multiple splice variants. There are multiple pseudogenes for this gene on chromosomes 1, 6, 7, 8, 9, and 13.

Application Notes

Optimal dilution of the Beta Tubulin antibody should be determined by the researcher.

Immunogen

Amino acids EQFTAMFRRKAFLHWYTGEGMDEMEFTEAE were used as the immunogen for the Beta Tubulin antibody.

Storage

After reconstitution, the Beta Tubulin antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

