


Beta Tubulin Antibody / TUBB3 [clone TUBB3/3732] (V7659)


Catalog No.	Formulation	Size
V7659-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7659-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7659SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	TUBB3/3732
Purity	Protein G affinity chromatography
UniProt	Q13509
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 1-2ug/ml
Limitations	This Beta Tubulin antibody is available for research use only.




Western blot testing of human U-87 cel lysate with Beta Tubulin antibody (clone TUBB3/3732). Predicted molecular weight: ~50 kDa.



IHC staining of FFPE human bladder carcinoma with Beta Tubulin antibody (TUBB3/3732). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Beta Tubulin antibody (clone TUBB3/3732). These results demonstrate the foremost specificity of the TUBB3/3732 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Description

This gene encodes a class III member of the beta tubulin protein family. Beta tubulins are one of two core protein families (alpha and beta tubulins) that heterodimerize and assemble to form microtubules. This protein is primarily expressed in neurons and may be involved in neurogenesis and axon guidance and maintenance. Mutations in this gene are the cause of congenital fibrosis of the extraocular muscles type 3. In adults, tubulin beta 3 (TUBB3) is primarily expressed in neurons and is commonly used as a neuronal marker. It plays an important role in neuronal cell proliferation and differentiation.

Application Notes

Optimal dilution of the Beta Tubulin antibody should be determined by the researcher.

Immunogen

Amino acids 437-450 were used as the immunogen for the Beta Tubulin antibody.

Storage

Store the Beta Tubulin antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).