


BMP15 Antibody / Bone morphogenetic protein 15 (RQ4880)


Catalog No.	Formulation	Size
RQ4880	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	O95972
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This BMP15 antibody is available for research use only.




IHC staining of FFPE human thyroid cancer with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.




IHC staining of FFPE human ovarian cancer with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.




IHC staining of FFPE human stomach cancer with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.




IHC staining of FFPE human lung cancer with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.




Western blot testing of 1) human HeLa, 2) human placenta, 3) human Caco-2, 4) rat lung and 5 mouse lung lysate with BPM15 antibody at 0.5ug/ml. Routinely observed molecular weight: ~50 kDa pro form and 16-17 kDa mature form.




IHC staining of FFPE human intestinal cancer with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.




IHC staining of FFPE human tonsil with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.



IHC staining of FFPE human placenta with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.



IHC staining of FFPE mouse lung with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.



IHC staining of FFPE rat lung with BMP15 antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.

Description

Bone morphogenetic protein 15 is a protein that in humans is encoded by the BMP15 gene. It is mapped to Xp11.22. It's mainly involved in folliculogenesis. The protein encoded by this gene is a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily. The transforming growth factor-beta superfamily includes large families of growth and differentiation factors. It is thought that this protein may be involved in oocyte maturation and follicular development as a homodimer or by forming heterodimers with a related protein, Gdf9. Defects in this gene are the cause of ovarian dysgenesis 2.

Application Notes

Optimal dilution of the BMP15 antibody should be determined by the researcher.

Immunogen

Amino acids KGTCLRVLRLDGLNSPNHAIQNLINQLVDQ from the human protein were used as the immunogen for the BMP15 antibody.

Storage

After reconstitution, the BMP15 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.