


## CA1 Antibody / Carbonic Anhydrase I [clone 2B5] (RQ6240)


| Catalog No. | Formulation                                           | Size   |
|-------------|-------------------------------------------------------|--------|
| RQ6240      | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

**Bulk quote request**


|                           |                                                                   |
|---------------------------|-------------------------------------------------------------------|
| <b>Availability</b>       | 1-3 business days                                                 |
| <b>Species Reactivity</b> | Human, Mouse, Rat                                                 |
| <b>Format</b>             | Antigen affinity purified                                         |
| <b>Clonality</b>          | Monoclonal (mouse origin)                                         |
| <b>Isotype</b>            | Mouse IgG1                                                        |
| <b>Clone Name</b>         | 2B5                                                               |
| <b>Purity</b>             | Affinity purified                                                 |
| <b>Buffer</b>             | Lyophilized from 1X PBS with 2% Trehalose                         |
| <b>UniProt</b>            | P00915                                                            |
| <b>Localization</b>       | Cytoplasmic                                                       |
| <b>Applications</b>       | Western Blot : 1-2ug/ml<br>Immunohistochemistry (FFPE) : 2-5ug/ml |
| <b>Limitations</b>        | This CA1 antibody is available for research use only.             |




IHC staining of FFPE human gastric cancer with CA1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human lung cancer with CA1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human tonsil with CA1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human colon cancer with CA1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human liver cancer with CA1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human pancreatic cancer with CA1 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) human K562, 2) rat spleen and 3) mouse spleen lysate with CA1 antibody. Predicted molecular weight: ~29 kDa.

## Description

Carbonic anhydrase 1 is an enzyme that in humans is encoded by the CA1 gene. It is a member of the Carbonic anhydrase. The CA1 gene is mapped to 8q22. CAI has got about 260 amino acids. This protein is highly expressed in erythrocytes. As catalysts of the reversible hydration of carbon dioxide, CAI participates in a variety of biologic processes like respiration, calcification, acid-base balance etc.

## Application Notes

Optimal dilution of the CA1 antibody should be determined by the researcher.

## Immunogen

A human recombinant partial protein (amino acids D9-F261) was used as the immunogen for the CA1 antibody.

## Storage

After reconstitution, the CA1 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.