


Cadherin 17 Antibody / CDH17 [clone CDH17/2612] (V5638)


Catalog No.	Formulation	Size
V5638-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5638-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5638SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	CDH17/2612
Purity	Protein G affinity
UniProt	Q12864
Localization	Cell membrane, cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 1-2ug/ml
Limitations	This Cadherin 17 antibody is available for research use only.



Western blot testing of human small intestine tissue lysate with Cadherin 17 antibody (clone CDH17/2612). Predicted molecular weight ~92 kDa but may be observed at higher molecular weights due to glycosylation.



IHC staining of FFPE human colon tissue with Cadherin 17 antibody (clone CDH17/2612). Inset: PBS used in place of primary Ab (secondary Ab negative control).
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Cadherin 17 antibody (clone CDH17/2612) as confirmation of integrity and purity.

Description

Cadherin 17 (also known as LI Cadherin) is part of a family of Calcium-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. Cadherins each contain a large extracellular domain at the amino terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short carboxy terminal, intracellular domain interacts with a variety of cytoplasmic proteins, including beta-catenin, to regulate cadherin function. LI-cadherin (for liver-intestine-cadherin) expression is restricted to liver and intestine tissues and is specifically localized to the basolateral domain of hepatocytes and enterocytes.

Application Notes

Optimal dilution of the Cadherin 17 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 242-418 from human Cadherin 17 protein was used as the immunogen for the Cadherin 17 antibody.

Storage

Aliquot the Cadherin 17 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.