


## CASP7 Antibody / Caspase 7 [clone CASP7/9054] (V5565)


| Catalog No.    | Formulation                                                             | Size   |
|----------------|-------------------------------------------------------------------------|--------|
| V5565-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V5565-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug  |
| V5565SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                          | 100 ug |

[Bulk quote request](#)

|                           |                                                         |
|---------------------------|---------------------------------------------------------|
| <b>Availability</b>       | 1-3 business days                                       |
| <b>Species Reactivity</b> | Human                                                   |
| <b>Format</b>             | Purified                                                |
| <b>Clonality</b>          | Monoclonal (mouse origin)                               |
| <b>Isotype</b>            | Mouse IgG2b, kappa                                      |
| <b>Clone Name</b>         | CASP7/9054                                              |
| <b>Purity</b>             | Protein A/G affinity                                    |
| <b>UniProt</b>            | P55210                                                  |
| <b>Localization</b>       | Cytoplasm                                               |
| <b>Applications</b>       | Immunohistochemistry (FFPE) : 1-2ug/ml                  |
| <b>Limitations</b>        | This CASP7 antibody is available for research use only. |



IHC staining of FFPE human kidney tissue with CASP7 antibody (clone CASP7/9054). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CASP7 antibody (clone CASP7/9051) as confirmation of integrity and purity.

## Description

A unique family of Cysteine proteases has been described that differs in sequence, structure and substrate specificity from any previously described protease family. This family, CED-3/caspase-1, is comprised of caspase-1, caspase-2, caspase-3, caspase-4, caspase-6, caspase-7 (also designated Mch3, ICE-LAP3 or CMH-1), caspase-9 and caspase-10. CED-3/caspase-1 family members function as key components of the apoptotic machinery and act to destroy specific target proteins which are critical to cellular longevity. Poly(ADP-ribose) polymerase plays an integral role in surveying for DNA mutations and double strand breaks. Caspase-3, caspase-7 and caspase-9, but not caspase-1, have been shown to cleave the nuclear protein PARP into an apoptotic fragment. Caspase-6, but not caspase-3, has been shown to cleave the nuclear lamins which are critical to maintaining the integrity of the nuclear envelope and cellular morphology. Caspase-10 has been shown to activate caspase-3 and caspase-7 in response to apoptotic stimuli.

## Application Notes

Optimal dilution of the CASP7 antibody should be determined by the researcher.

## Immunogen

A recombinant fragment (within amino acids 1-200) of human CASP7 protein was used as the immunogen for the CASP7 antibody.

## Storage

Aliquot the CASP7 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.