


CD10 Antibody / Neutral endopeptidase 24.11 / Neprilysin [clone MME/8658R] (V5014)

Catalog No.	Formulation	Size
V5014-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5014-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5014SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	MME/8658R
Purity	Protein A/G affinity
UniProt	P08473
Localization	Cell surface, Cytoplasm
Applications	Western Blot : 2-4ug/ml
Limitations	This CD10 antibody is available for research use only.



Western blot testing of human kidney tissue lysate with CD10 antibody. Routinely visualized at ~100 kDa.



Western blot testing of human kidney tissue lysate with CD10 antibody. Routinely visualized at ~100 kDa.



SDS-PAGE analysis of purified, BSA-free CD10 antibody (clone MME/8658R) as confirmation of integrity and purity.

Description

Recognizes a 100kDa glycoprotein, identified as CD10, also known as Common Acute Lymphocytic Leukemia Antigen (CALLA). It is a cell surface enzyme with neutral metalloendopeptidase activity, which inactivates a variety of biologically active peptides. CD10 is expressed on the cells of lymphoblastic, Burkitt's, and follicular germinal center lymphomas, and on cells from patients with chronic myelocytic leukemia (CML). It is also expressed on the surface of normal early lymphoid progenitor cells, immature B cells within adult bone marrow and germinal center B cells within lymphoid tissue. CD10 is also present on breast myoepithelial cells, bile canaliculi, fibroblasts, with especially high expression on the brush border of kidney and gut epithelial cells.

Application Notes

Optimal dilution of the CD10 antibody should be determined by the researcher.

Immunogen

A recombinant human CD10 protein fragment (within amino acids 1-500) was used as the immunogen for the CD10 antibody.

Storage

Aliquot the CD10 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.