


CD105 Antibody [clone CDLA105-1] (V8311)

| Catalog No. | Formulation | Size |
|----------------|--|--------|
| V8311-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V8311-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V8311SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Bulk quote request

| Availability | 1-3 business days |
|--------------------|---|
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG2b, kappa |
| Clone Name | CDLA105-1 |
| Purity | Protein G affinity chromatography |
| UniProt | P17813 |
| Localization | Cell surface, cytoplasmic |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml |
| Limitations | This CD105 antibody is available for research use only. |



IHC staining of FFPE human tonsil with CD105 antibody (clone CDLA105-1). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.

Description

Vascular endothelium glycoprotein that plays an important role in the regulation of angiogenesis. Required for normal structure and integrity of adult vasculature. Regulates the migration of vascular endothelial cells. Required for normal

extraembryonic angiogenesis and for embryonic heart development (By similarity). May regulate endothelial cell shape changes in response to blood flow, which drive vascular remodeling and establishment of normal vascular morphology during angiogenesis (By similarity). May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors. [UniProt]

Application Notes

The stated application concentrations are suggested starting points. Titration of the CD105 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Amino acids 74-251 were used as the immunogen for the CD105 antibody.

Storage

Store the CD105 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).