

## CD133 Antibody / PROM1 [clone 6H10-F1-C11] (F54026)


| Catalog No.  | Formulation   | Size   |
|--------------|---|--------|
| F54026-0.1ML | In PBS with 50% glycerol, 0.1mg/ml BSA and 0.02% sodium azide | 0.1 ml |

**Bulk quote request**

|                           |   |
|---------------------------|---|
| <b>Availability</b>       | 1-3 business days                                       |
| <b>Species Reactivity</b> | Human   |
| <b>Format</b>             | Purified  |
| <b>Clonality</b>          | Monoclonal (mouse origin)                               |
| <b>Isotype</b>            | Mouse IgG2a   |
| <b>Clone Name</b>         | 6H10-F1-C11   |
| <b>Purity</b>             | Protein G affinity                                      |
| <b>UniProt</b>            | O43490  |
| <b>Applications</b>       | Western Blot : 1:1000                                   |
| <b>Limitations</b>        | This CD133 antibody is available for research use only. |



Western blot testing of human CaCo2 cell lysate with CD133 antibody at 1:1000. Predicted molecular weight: ~97 kDa (unmodified), ~130 kDa (glycosylated).



Western blot testing of A) untransfected and B) transfected CHO-K1 cells with CD133 antibody at 1:1000. Predicted molecular weight: ~97 kDa (unmodified), ~130 kDa (glycosylated).

## Description

CD133/PROM1/Prominin-1 may play a role in cell differentiation, proliferation and apoptosis. Binds cholesterol in cholesterol-containing plasma membrane microdomains and may play a role in the organization of the apical plasma membrane in epithelial cells. During early retinal development acts as a key regulator of disk morphogenesis. Involved in regulation of MAPK and Akt signaling pathways. In neuroblastoma cells suppresses cell differentiation such as neurite outgrowth in a RET-dependent manner. [UniProt]

## Application Notes

The stated application concentrations are suggested starting points. Titration of the CD133 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A human recombinant partial protein was used as the immunogen for this CD133 antibody.

## Storage

Store the CD133 antibody at -20oC.