


## CD23 Antibody [clone FCER2/8234R] (V4599)

| Catalog No.    | Formulation                                                             | Size   |
|----------------|-------------------------------------------------------------------------|--------|
| V4599-100UG    | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V4599-20UG     | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug  |
| V4599SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free                          | 100 ug |


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

|                    |                                                         |
|--------------------|---------------------------------------------------------|
| Availability       | 1-3 business days                                       |
| Species Reactivity | Human                                                   |
| Format             | Purified                                                |
| Clonality          | Recombinant Rabbit Monoclonal                           |
| Isotype            | Rabbit IgG, kappa                                       |
| Clone Name         | FCER2/8234R                                             |
| Purity             | Protein A/G affinity                                    |
| UniProt            | P06734                                                  |
| Localization       | Cell Surface                                            |
| Applications       | Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT |
| Limitations        | This CD23 antibody is available for research use only.  |



IHC staining of FFPE human tonsil tissue with CD23 antibody (clone FCER2/8234R). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CD23 antibody (clone FCER2/8234R) as confirmation of integrity and purity.

## Description

CD23 (FCE2) is a type II integral membrane glycoprotein that is expressed on mature B cells, monocytes, eosinophils, platelets and dendritic cells. CD23 is a low affinity IgE receptor that mediates IgE-dependent cytotoxicity and phagocytosis by macrophages and eosinophils. CD23 associates as an oligomer where cooperative binding of at least two lectin domains is required for high affinity IgE binding to CD23. It may play a role in antigen presentation by B cells by interacting with CD40. CD23 has been shown to be associated with the Fyn tyrosine kinase. The truncated molecule can be secreted, then function as a potent mitogenic growth factor. CD23 is expressed on a subpopulation of peripheral blood cells, B-lymphocytes and on EBV transformed B lymphoblastoid cell lines. CD23 is also detected in neoplastic cells from cases of B cell chronic lymphocytic leukemia and some cases on centroblastic/centrocytic lymphoma.

## Application Notes

Optimal dilution of the CD23 antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein sequence (within amino acids 200-300) from the human protein was used as the immunogen for the CD23 antibody.

## Storage

Aliquot the CD23 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.