


CD71 Antibody / Transferrin Receptor [clone 66IG10] (V2261CF488)


Catalog No.	Formulation	Size
V2261CF488-100T	500 ul at 0.1 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 Tests

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	CF488 Conjugate
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	66IG10
Purity	Protein G affinity chromatography
UniProt	P02786
Localization	Cell surface
Applications	Flow Cytometry : 1-2ug/million cells Immunofluorescence : 1-3ug/ml
Limitations	This CD71 antibody is available for research use only.



Flow cytometry testing of K562 cells with AF488 conjugated CD71 antibody (green, clone 66IG10), cells alone (black) and isotype control (gray).



FACS testing of human Jurkat cells with CF488 conjugated Transferrin receptor antibody (blue) and isotype control (red).

Description

It recognizes a ~90-95kDa protein which is identified as cell surface transferrin receptor (CD71), a disulfide-bonded homodimeric glycoprotein of 180-190kDa. This MAb is highly specific to CD71 and shows no cross-reaction with other related proteins. Its epitope is localized in the extracellular domain of CD71. Ligand for transferrin receptor is the serum iron transport protein, transferrin. This receptor is broadly distributed in carcinomas, sarcomas, leukemias, and lymphomas. CD71/Transferrin receptor has been reported to be associated with cell proliferation in both normal and neoplastic tissues and useful in predicting clinical behavior or response to therapy in a number of malignancies including breast cancer.

Application Notes

Optimal dilution of the CD71 antibody should be determined by the researcher.

Immunogen

Recombinant human protein was used as the immunogen for this Transferrin receptor / CD71 antibody.

Storage

Store the CD71 antibody at 2-8°C, protected from light.