


## CDC5L Antibody / Cell division cycle 5-like protein [clone PCR- CDC5L-2C6] (V5121)


Catalog No.	Formulation	Size
V5121-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5121-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5121SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2a, kappa
<b>Clone Name</b>	PCR-CDC5L-2C6
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	Q99459
<b>Localization</b>	Nucleus, Cytoplasm
<b>Applications</b>	Flow Cytometry : 1-2ug/million cells Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
<b>Limitations</b>	This CDC5L antibody is available for research use only.



Flow cytometry testing of human HeLa cells with CDC5L antibody (clone PCR-CDC5L-2C6) followed by goat anti-mouse IgG-CF488 (blue); isotype control (red).



IHC staining of FFPE human colon carcinoma tissue with CDC5L antibody (clone PCRP-CDC5L-2C6). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Analysis of a HuProt(TM) microarray containing more than 19,000 full-length human proteins using CDC5L antibody (clone PCRP-CDC5L-2C6) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD&#39;s) above the mean value of all signals generated on that array. If targets on HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD&#39;s) between the Z-score. S-score therefore represents the relative target specificity of a mAb to its intended target. A mAb is considered to specific to its intended target, if the mAb has an S-score of at least 2.5. For example, if a mAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that mAb to protein X is equal to 29.

## Description

Cdc5L (cell division cycle 5-like protein, pombe Cdc5-related protein) is a DNA-binding protein encoded by the human gene CDC5L. Cdc5L contains two HTH Myb-type DNA-binding domains and may shuttle between cytoplasm and nucleus. It is involved in cell cycle control and may act as a transcription activator. Cdc5L is a spliceosomal protein that is highly conserved across species. It belongs to the group of proteins that comprise the core of spliceosomal complexes and are essential for pre-mRNA splicing. Cdc5L is involved in the second catalytic step of pre-mRNA splicing, which involves cleavage at the 3' splice site and the ligation of the exons. This process releases the intact intron lariat. A chromosomal aberration involving Cdc5L is found in multicystic renal dysplasia. This aberration is caused by a translocation (t 6;19,p21;q13.1) with USF-2.

## Application Notes

Optimal dilution of the CDC5L antibody should be determined by the researcher.

## Immunogen

A recombinant partial protein sequence (within amino acids 1-148) from the human protein was used as the immunogen for the CDC5L antibody.

## Storage

Aliquot the CDC5L antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

