


CDH1 Antibody [clone 813CT11.1.3] (F52228)


| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F52228-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F52228-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

Bulk quote request

| Availability | 1-3 business days |
|--------------------|--|
| Species Reactivity | Human, Mouse |
| Format | Purified |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG1, k |
| Clone Name | 813CT11.1.3 |
| Purity | Purified |
| UniProt | P12830 |
| Localization | Cytoplasmic, membranous |
| Applications | IHC (Paraffin): 1:25 Western Blot: 1:1000 |
| Limitations | This CDH1 antibody is available for research use only. |



Immunohistochemical analysis of paraffin-embedded human stomach using CDH1 antibody at 1:25 dilution.



CDH1 antibody western blot analysis in (1) human WiDr cell line and (2) mouse stomach tissue lysate. Expected molecular weight: 135 kDa (precursor), 80-120 kDa (mature, depending on gylcosylation level).

Description

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells. Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7. E-Cad/CTF2 promotes non-amyloidogenic degradation of Abeta precursors. Has a strong inhibitory effect on APP C99 and C83 production.

Application Notes

Titration of the CDH1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Purified His-tagged protein was used to produced this monoclonal CDH1 antibody.

Storage

Aliquot the CDH1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.