


CK1 Antibody / Cytokeratin 1 / Keratin 1 [clone KRT1/1840] (V7908)


Catalog No.	Formulation	Size
V7908-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7908-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7908SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	KRT1/1840
Purity	Protein G affinity chromatography
UniProt	P04264
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 2-4ug/ml
Limitations	This CK1 antibody is available for research use only.



IHC staining of FFPE human cervical carcinoma with CK1 antibody (clone KRT1/1840).
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human skin with CK1 antibody (clone KRT1/1840). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free CK1 antibody (clone KRT1/1840) as confirmation of integrity and purity.

Description

Keratins are a family of intermediate filament proteins that assemble into filaments through forming heterodimers of one type I keratin (keratins 9 to 23) and one type II keratin (keratins 1 to 8). Keratins demonstrate tissue and differentiation specific expression profiles. Cytokeratin 1 is a hetero-tetramer of two type I as well as two type II keratins, keratin-10 (KRT10). Keratin 1 is differentiation specific keratin that is one of the predominant keratins in suprabasal keratinocytes in stratified epithelia. Mutations in keratin 1 cause epidermolytic hyperkeratosis.

Application Notes

Optimal dilution of the CK1 antibody should be determined by the researcher.

Immunogen

A recombinant full-length human Cytokeratin 1 protein was used as the immunogen for this CK1 antibody.

Storage

Store the CK1 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).