


Cofilin Antibody (R31465)


Catalog No.	Formulation	Size
R31465	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
Gene ID	1072
Localization	Cytoplasmic, membrane
Applications	Western Blot : 0.5-1ug/ml IHC (FFPE) : 0.5-1ug/ml IF/ICC (FFPE) : 2-4ug/ml ELISA : 0.1-0.5ug/ml (human protein tested); request BSA free format for coating
Limitations	This Cofilin antibody is available for research use only.




Western blot testing of 1) human HeLa, 2) monkey COS-7, human 3) HEK293, 4) U-87 MG, 5) HepG2, 6) PC-3, 7) ThP-1 and 8) A549 cell lysate with Cofilin antibody. Predicted molecular weight ~19 kDa.




Western blot testing 1) rat heart, 2) rat spleen, 3) rat lung, 4) rat C6, 5) mouse spleen and 6) mouse HEPA1-6 lysate with Cofilin antibody. Predicted molecular weight ~19 kDa.




IHC-P: Cofilin antibody testing of human breast cancer tissue. HIER: steam section in pH6 citrate buffer for 20 min.




IHC-P: Cofilin antibody testing of human intestine cancer tissue. HIER: steam section in pH6 citrate buffer for 20 min.




IHC-P testing of mouse lung tissue. HIER: steam section in pH6 citrate buffer for 20 min.




IHC-P testing of rat lung tissue. HIER: steam section in pH6 citrate buffer for 20 min.



IF/ICC staining of FFPE human U-2 OS cells with Cofilin antibody (green) at 2ug/ml. HIER: steam section in pH6 citrate buffer for 20 min.



IHC staining of frozen human placenta with Cofilin antibody.



Flow cytometry testing of human U-2 OS cells with Cofilin antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Cofilin antibody.

Description

Cofilin is a ubiquitous actin-binding factor required for the reorganization of actin filaments. The protein is known to sever actin filaments by creating more positive ends on filament fragments, it can cause depolymerization at the minus end of filaments, thereby preventing their reassembly. As a long-lasting in vivo effect, Cofilin recycles older ADP-F-actin, helping cell to maintain ATP-G-actin pool for sustained motility. pH, phosphorylation and phosphoinositides regulate cofilins binding and associating activity with actin.

Application Notes

The stated application concentrations are suggested starting amounts. Titration of the Cofilin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Human partial recombinant protein (AA 2-166) was used as the immunogen for this Cofilin antibody.

Storage

After reconstitution, the Cofilin antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.