


COL4A Antibody / Collagen IV [clone rCOL4/4742] (V8646)

| Catalog No. | Formulation | Size |
|----------------|--|--------|
| V8646-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V8646-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V8646SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |


Recombinant MOUSE MONOCLONAL

Bulk quote request


| Availability | 1-3 business days |
|--------------------|--|
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Recombinant Mouse Monoclonal |
| Isotype | Mouse IgG2a, kappa |
| Clone Name | rCOL4/4742 |
| Purity | Protein G affinity chromatography |
| UniProt | P02462 |
| Applications | Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT |
| Limitations | This COL4A antibody is available for research use only. |



IHC staining of FFPE human kidney with recombinant COL4A antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human kidney with recombinant COL4A antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free COL4A antibody as confirmation of integrity and purity.

Description

Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers.

Application Notes

Optimal dilution of the COL4A antibody should be determined by the researcher.

Immunogen

Purified human placental extract was used as the immunogen for the COL4A antibody.

Storage

Store the COL4A antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).