


Cxcr3 Antibody (RQ7070)


Catalog No.	Formulation	Size
RQ7070	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	O88410
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This Cxcr3 antibody is available for research use only.



IHC staining of FFPE mouse intestinal lymph node tissue with Cxcr3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE rat lymph node tissue with Cxcr3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) rat C6 and 2) mouse NIH 3T3 cell lysate with Cxcr3 antibody.
Predicted molecular weight ~41 kDa.

Description

Chemokine receptor CXCR3 is a G α protein-coupled receptor in the CXC chemokine receptor family. Other names for CXCR3 are G protein-coupled receptor 9 (GPR9) and CD183. It is mapped to Xq13.1. CXCR3 is expressed on malignant B cells from chronic lymphoproliferative disorders, particularly in patients with CLL, and represents a fully functional receptor involved in chemotaxis of malignant B lymphocytes. It is found that in the absence of known etiologic agents, CXCR3 represents a novel target for therapeutic interference early in type 1 diabetes. CXCR3 signaling is associated with MG pathogenesis and proposed that CXCR3 may serve as novel drug targets to treat MG. CXCR3A and CXCR3B are involved in the chemotactic and vascular effects of CXCL4L1.

Application Notes

Optimal dilution of the Cxcr3 antibody should be determined by the researcher.

Immunogen

Recombinant mouse protein (amino acids M1-E326) was used as the immunogen for the Cxcr3 antibody.

Storage

After reconstitution, the Cxcr3 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.