


# Cytokeratin 1 Antibody / CK1 / Keratin 1 [clone LHK1] (V7907)


Catalog No.	Formulation	Size
V7907-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7907-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7907SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

# **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	LHK1
Purity	Protein G affinity chromatography
UniProt	P04264
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE): 0.1-0.2ug/ml
Limitations	This Cytokeratin 1 antibody is available for research use only.



IHC staining of FFPE human cervix with Cytokeratin 1 antibody (clone LHK1). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human skin with Cytokeratin 1 antibody (clone LHK1). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Cytokeratin 1 antibody (clone LHK1) as confirmation of integrity and purity.

## **Description**

Keratins are a family of intermediate filament proteins that assemble into filaments through forming heterodimers of one type I keratin (keratins 9 to 23) and one type II keratin (keratins 1 to 8). Keratins demonstrate tissue and differentiation specific expression profiles. Cytokeratin 1 is a hetero-tetramer of two type I as well as two type II keratins, keratin-10 (KRT10). Keratin 1 is differentiation specific keratin that is one of the predominant keratins in suprabasal keratinocytes in stratified epithelia. Mutations in keratin 1 cause epidermolytic hyperkeratosis.

### **Application Notes**

Optimal dilution of the Cytokeratin 1 antibody should be determined by the researcher.

#### **Immunogen**

Amino acids VRFVSTSYSGVTR from the C-terminal portion of the protein was used as the immunogen for this Cytokeratin 1 antibody.

#### **Storage**

Store the Cytokeratin 1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).