



# DcR3 Antibody / TNFRSF6B (F54535)

Catalog No.	Formulation	Size
F54535-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54535-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml


#### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity purified
UniProt	O95407
Applications	Western Blot : 1:500-1:2000 Flow Cytometry : 1:25 (1x10e6 cells) Immunohistochemistry (FFPE) : 1:25
Limitations	This DcR3 antibody is available for research use only.






Western blot testing of human 1) HeLa, 2) HCT116, 3) PC-3, 4) SW480 and 5) HUVEC cell lysate with DcR3 antibody. Predicted molecular weight ~33 kDa.



IHC testing of FFPE human colon carcinoma tissue with DcR3 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Flow cytometry testing of human HEK293 cells with DcR3 antibody; Blue=isotype control, Green= DcR3 antibody.

## **Description**

This gene belongs to the tumor necrosis factor receptor superfamily. The encoded protein is postulated to play a regulatory role in suppressing FasL- and LIGHT-mediated cell death. It acts as a decoy receptor that competes with death receptors for ligand binding. Overexpression of this gene has been noted in gastrointestinal tract tumors, and it is located in a gene-rich cluster on chromosome 20, with other potentially tumor-related genes. Two transcript variants encoding the same isoform, but differing in the 5' UTR, have been observed for this gene.

## **Application Notes**

The stated application concentrations are suggested starting points. Titration of the DcR3 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A portion of amino acids 22-48 from the human protein was used as the immunogen for the DcR3 antibody.

#### **Storage**

Aliquot the DcR3 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.