


## Diphosphomevalonate decarboxylase Antibody / MVD (RQ8521)


Catalog No.	Formulation	Size
RQ8521	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 days
<b>Species Reactivity</b>	Human, Mouse
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity purified
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose
<b>UniProt</b>	P53602
<b>Localization</b>	Cytoplasm
<b>Applications</b>	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells ELISA : 0.1-0.5ug/ml
<b>Limitations</b>	This Diphosphomevalonate decarboxylase antibody is available for research use only.



IHC staining of FFPE mouse ovary tissue with Diphosphomevalonate decarboxylase antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) HepG2, 2) MCF7, 3) K562 and 4) Caco-2 cell lysate with Diphosphomevalonate decarboxylase antibody. Predicted molecular weight ~43 kDa.



Flow cytometry testing of fixed and permeabilized human MCF7 cells with Diphosphomevalonate decarboxylase antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Diphosphomevalonate decarboxylase antibody.

## Description

The enzyme mevalonate pyrophosphate decarboxylase (MVD; EC 4.1.1.33) catalyzes the conversion of mevalonate pyrophosphate into isopentenyl pyrophosphate. This unusual enzyme decarboxylates and dehydrates its substrate while hydrolyzing ATP. As a unique enzyme in one of the early steps in cholesterol biosynthesis, MVD may be a useful target for drugs aimed at lowering serum cholesterol levels. This gene is mapped to chromosome 16q24.3 based on an alignment of the MVD sequence.

## Application Notes

Optimal dilution of the Diphosphomevalonate decarboxylase antibody should be determined by the researcher.

## Immunogen

An E.coli-derived human recombinant protein (amino acids E62-Q374) was used as the immunogen for the Diphosphomevalonate decarboxylase antibody.

## Storage

After reconstitution, the Diphosphomevalonate decarboxylase antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.