


Dopamine Beta-Hydroxylase Antibody / DBH [clone DBH/7222] (V5312)

Catalog No.	Formulation	Size
V5312-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5312-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5312SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2, kappa
Clone Name	DBH/7222
Purity	Protein A/G affinity
UniProt	P09172
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Dopamine Beta-Hydroxylase antibody is available for research use only.



IHC staining of FFPE human adrenal gland tissue with Dopamine Beta-Hydroxylase antibody (clone DBH/7222). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

Dopamine beta-hydroxylase (DBH) catalyzes the conversion of dopamine to noradrenaline in the biosynthesis of catecholamines. DBH is selectively expressed in noradrenergic and adrenergic neurons, as well as in neuroendocrine cells, and it serves as a specific protein marker for noradrenergic processes. The active form of DBH is a homotetramer, which is found in the lumen of synaptic vesicles of corresponding nerve cells, where it localizes to both the membrane

and cytosol. DBH is induced by nerve growth factor and Insulin growth factor-1 and is regulated by intracellular second messengers protein kinase A, cyclic AMP, diacyl glycerol and Ca²⁺. Expression of DBH is transcriptionally mediated by Sp1, CREB and AP-1 proteins including c-Fos, c-Jun and JunD.

Application Notes

Optimal dilution of the Dopamine Beta-Hydroxylase antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 200-400) from the human protein was used as the immunogen for the Dopamine Beta-Hydroxylase antibody.

Storage

Aliquot the Dopamine Beta-Hydroxylase antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.