


Double Stranded DNA Antibody / dsDNA [clone rDSD/8266] (V5319)

Catalog No.	Formulation	Size
V5319-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5319-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5319SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG2b, kappa
Clone Name	rDSD/8266
Purity	Protein A/G affinity
UniProt	Not Applicable
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This Double Stranded DNA antibody is available for research use only.



IHC staining of FFPE human ovarian cancer tissue with Double Stranded DNA antibody (clone rDSD/8266). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Double Stranded DNA antibody (clone rDSD/8266) as confirmation of integrity and purity.

Description

This mAb recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This mAb produces a homogeneous staining pattern in the nucleus of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3',5'-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.

Application Notes

Optimal dilution of the Double Stranded DNA antibody should be determined by the researcher.

Immunogen

Burkitt's cell nuclei were used as the immunogen for the Double Stranded DNA antibody.

Storage

Aliquot the Double Stranded DNA antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.