


dsDNA Antibody [clone DRNC1-1] (V7189)


Catalog No.	Formulation	Size
V7189-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7189-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7189SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7189IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG3, kappa
Clone Name	DRNC1-1
Purity	Protein G affinity chromatography
UniProt	Not Known
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT (1)
Limitations	This dsDNA antibody is available for research use only.



IHC testing of FFPE human tonsil with dsDNA antibody (clone DRNC1-1). FFPE testing requires sections to be boiled in pH 9 10mM Tris with 1mM EDTA for 10-20 minutes, followed by cooling at RT for 20 minutes, prior to staining.



IHC testing of FFPE human colon carcinoma with dsDNA antibody (clone DRNC1-1). FFPE testing requires sections to be boiled in pH 9 10mM Tris with 1mM EDTA for 10-20 minutes, followed by cooling at RT for 20 minutes, prior to staining.

Description

Deoxyribonucleic acid (DNA) is a nucleic acid that stores long-term information regarding the development and function of all known living organisms. DNA consists of two long nucleotide polymers, which are composed of four bases, namely adenine, thymine, guanine and cytosine, all of which are flanked by a phosphate-deoxyribose backbone. This mAb can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells.

Application Notes

Titration of the dsDNA antibody may be required for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Cell nuclei were used as the immunogen for the dsDNA antibody.

Storage

Store the dsDNA antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).