


## Eotaxin Antibody (R31494)

Catalog No.	Formulation	Size
R31494	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Mouse
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide
<b>Gene ID</b>	20292
<b>Applications</b>	Western Blot : 0.5-1ug/ml
<b>Limitations</b>	This Eotaxin antibody is available for research use only.



Western blot testing of Eotaxin antibody (0.5ug/ml) and Lane 1: mouse lung; 2: (m) intestine; 3: (m) HEPA lysate

## Description

CCL11, also known as Eotaxin, is a potent inducer of eosinophil chemotaxis and is considered as a selective ligand of the CC chemokine receptor 3 (CCR3), which is expressed on eosinophils, basophils, and Th2 lymphocytes. The gene maps to chromosome 17 and is expressed constitutively at high levels in small intestine and colon, and at lower levels in various other tissues. CCL11/Eotaxin is a small cytokine belonging to the CC chemokine family. It is assumed to be involved in eosinophilic inflammatory diseases such as atopic dermatitis, allergic rhinitis, asthma and parasitic infections. The effects of Eotaxin are mediated by its binding to a G-protein-linked receptor known as a chemokine receptor.

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the Eotaxin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## **Immunogen**

Mouse partial recombinant protein (AA 24-97) was used as the immunogen for this Eotaxin antibody.

## **Storage**

After reconstitution, the Eotaxin antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.