

# Eph Receptor B2 Antibody [EphB2] (F50585)

Catalog No.	Formulation	Size
F50585-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F50585-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Purified
UniProt	P29323
Applications	IHC (Paraffin): 1:10-1:100
Limitations	This EphB2 antibody is available for research use only.



EphB2 antibody analysis in formalin fixed and paraffin embedded human skeletal muscle

## **Description**

Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The ligand-activated form of EphB2, which belongs to the Tyr family of protein kinases, interacts with multiple proteins, including GTPase-activating protein (RASGAP) through its SH2 domain. It binds RASGAP through the juxtamembrane tyrosines residues, and also interacts with PRKCABP and

GRIP1 This type I membrane protein is expressed in brain, heart, lung, kidney, placenta, pancreas, liver and skeletal muscle. It is preferentially expressed in fetal brain. This protein contains putatively 2 fibronectin type III domains and 1 sterile alpha motif (SAM) domain.

## **Application Notes**

Titration of the EphB2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

#### **Immunogen**

A portion of amino acids 103-133 from the human protein was used as the immunogen for this EphB2 antibody.

#### **Storage**

Aliquot the EphB2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.