



FAP Antibody / Fibroblast Activation Protein Alpha [clone FAP/4854] (V9498)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V9498-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V9498-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V9498SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

[Bulk quote request](#)

| | |
|---------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Monoclonal (mouse origin) |
| Isotype | Mouse IgG2c, kappa |
| Clone Name | FAP/4854 |
| Purity | Protein A/G affinity |
| UniProt | Q12884 |
| Localization | Cell surface, Cell membrane |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml |
| Limitations | This FAP antibody is available for research use only. |





IHC staining of FFPE human placental tissue with FAP antibody (clone FAP/4854).
HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

FAP (fibroblast activation protein) is a cell surface glycoprotein and serine protease that is expressed primarily in fetal mesenchymal tissues and epithelial cancer fibroblasts. In cancer, FAP functions to promote cellular proliferation. In embryonic development, FAP functions to remodel developing tissues. FAP acts as an integral membrane gelatinase composed of N-glycosylated proteolytically inactive subunits. FAP expression on chondrocyte membranes is upregulated by the combination of the cytokines IL-1 and OSM and has been shown to increase in osteoarthritic patients. This expression is colocalized with MMP-1 and MMP-13 as well as CD44 (variants v3 and v7/8). Mice that lack all copies of the FAP gene have been found to be fertile and to have developmental defects or change in cancer susceptibility.

Application Notes

Optimal dilution of the FAP antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-200 was used as the immunogen for the FAP antibody.

Storage

Aliquot the FAP antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.