


Ferritin Light Chain Antibody [clone FTL/1388] (V3352)


Catalog No.	Formulation	Size
V3352-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3352-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3352SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2a, kappa
Clone Name	FTL/1388
Purity	Protein G affinity chromatography
UniProt	P02792
Localization	Cytoplasmic
Applications	Western Blot : 0.5-2ug/ml Immunohistochemistry (FFPE) : 0.1-0.2ug/ml for 30 min at RT
Limitations	This Ferritin Light Chain antibody is available for research use only.




IHC testing of FFPE human pancreas with Ferritin Light Chain antibody (clone FTL/1388). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min.




IHC testing of FFPE human pancreas with Ferritin Light Chain antibody (clone FTL/1388). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min.



SDS-PAGE Analysis of Purified, BSA-Free Ferritin Light Chain Antibody (clone FTL/1388). Confirmation of Integrity and Purity of the Antibody.




Western blot testing of human HeLa lysate with Ferritin Light Chain antibody. Predicted molecular weight: ~20 kDa.



Western blot testing of human kidney lysate with Ferritin Light Chain antibody. Predicted molecular weight: ~20 kDa.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using Ferritin Light Chain antibody. These results demonstrate the foremost specificity of the FTL/1388 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Description

Application Notes

Optimal dilution of the Ferritin Light Chain antibody should be determined by the researcher.

Immunogen

Amino acids 38-165 of human FTL were used as the immunogen for this Ferritin Light Chain antibody.

Storage

Store the Ferritin Light Chain antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

References (1)