


## GATA3 Antibody [clone GATA3/2441] (V8874)

Catalog No.	Formulation	Size
V8874-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V8874-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V8874SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Monoclonal (mouse origin)
<b>Isotype</b>	Mouse IgG2b, kappa
<b>Clone Name</b>	GATA3/2441
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	P23771
<b>Localization</b>	Nuclear
<b>Applications</b>	ELISA (order BSA-free Format For Coating) : Flow Cytometry : 1-2ug/million cells Western Blot : 1-2ug/ml
<b>Limitations</b>	This GATA3 antibody is available for research use only.

### Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using GATA3 antibody (clone GATA3/2441). These results demonstrate the foremost specificity of the GATA3/2441 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free GATA3 antibody (clone GATA3/2441) as confirmation of integrity and purity.

## Description

GATA-3 is a zinc finger transcription factor and plays an important role in promoting and directing cell proliferation, development, and differentiation in many tissues and cell types. GATA-3 expression is primarily seen in breast and urothelial carcinomas. Therefore, GATA3 antibody can be useful in the identification of unknown primary carcinoma when carcinomas of the breast or bladder are a possibility.

## Application Notes

Optimal dilution of the GATA3 antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 357-436 was used as the immunogen for the GATA3 antibody.

## Storage

Aliquot the GATA3 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.