


# GATA3 Antibody [clone GATA3/2446] (V7481)


Catalog No.	Formulation	Size
V7481-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7481-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7481SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

# **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	GATA3/2446
Purity	Protein G affinity chromatography
UniProt	P23771
Localization	Nuclear
Applications	ELISA (order BSA/sodium Azide-free Format For Coating) : Flow Cytometry : 1-2ug/10^6 cells Western Blot : 0.5-1ug/ml
Limitations	This GATA3 antibody is available for research use only.



IHC testing of FFPE human breast carcinoma with GATA3 antibody. Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min and allow to cool prior to testing.



SDS-PAGE analysis of purified, BSA-free GATA3 antibody as confirmation of integrity and purity.

## **Description**

GATA3 is a zinc finger transcription factor and plays an important role in promoting and directing cell proliferation, development, and differentiation in many tissues and cell types. GATA3 expression is primarily seen in breast and urothelial carcinomas. Therefore, GATA3 antibody can be useful in the identification of unknown primary carcinoma when carcinomas of the breast or bladder are a possibility.

### **Application Notes**

Optimal dilution of the GATA3 antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

#### **Immunogen**

A portion of amino acids 357-463 from the human protein were used as the immunogen for this GATA3 antibody.

#### **Storage**

Store the GATA3 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).