


HDAC2 Antibody (RQ6487)


Catalog No.	Formulation	Size
RQ6487	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q92769
Localization	Nuclear, cytoplasmic
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml
Limitations	This HDAC2 antibody is available for research use only.




IHC staining of FFPE human ovarian cancer tissue with HDAC2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human gastric cancer tissue with HDAC2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human cervical cancer tissue with HDAC2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer tissue with HDAC2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human endometrial cancer tissue with HDAC2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human gallbladder adenocarcinoma tissue with HDAC2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human lung cancer tissue with HDAC2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE mouse testis tissue with HDAC2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Immunofluorescent staining of FFPE human Caco-2 cells with HDAC2 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human K562, 2) human ThP-1, 3) human HL60, 4) human Caco-2, 5) human HepG2, 6) rat brain, 7) rat lung, 8) rat pancreas, 9) mouse brain and 10) mouse lung lysate with HDAC2 antibody. Predicted molecular weight: 55-60 kDa.

Description

Histone deacetylase 2 is an enzyme that in humans is encoded by the HDAC2 gene. This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes and are responsible for the deacetylation of lysine residues on the N-terminal region of the core histones (H2A, H2B, H3 and H4). This protein also forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus it plays an important role in transcriptional regulation, cell cycle progression and developmental events. Betz et al. (1998) performed PCR using HDAC2-specific primers to screen a somatic cell hybrid mapping panel. They mapped the HDAC2 gene to human chromosome 6q21, a region of the genome altered in some cancers, including retinoblastoma.

Application Notes

Optimal dilution of the HDAC2 antibody should be determined by the researcher.

Immunogen

An E. coli-derived human protein (amino acids E387-P488) was used as the immunogen for the HDAC2 antibody.

Storage

After reconstitution, the HDAC2 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

