


## HDAC9 Antibody (F41192)

Catalog No.	Formulation	Size
F41192-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F41192-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml


[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Predicted Reactivity	Mouse, Chicken
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Purified
UniProt	Q9UKV0
Applications	Western Blot : 1:1000 Immunoprecipitation : 1:100 Immunofluorescence : 1:1000 IHC (Paraffin) : 1:50-1:100
Limitations	This HDAC9 antibody is available for research use only.




HDAC9 antibody NSJ# F41192 and NSJ# F41193 were tested by WB and IP-WB using HeLa and HeLa-HDAC9 transfected cells. Both antibodies detect the protein in transfected cells but not non-transfected.


IP testing of HDAC9 Abs




HDAC9 antibody NSJ# F41192 and NSJ# F41193 can both immunoprecipitate the protein from HeLa-HDAC9 transfected cells. (Data kindly provided by Dr. Zhigang Yuan, H. Lee Moffitt Cancer Center and Research Institute)



Immunoblots for MITR (HDAC9), Mirk, MyoD and tubulin proteins are shown for cytoplasmic (Cyt) and nuclear (N) extracts from undifferentiated C2C12 myoblasts transfected with plasmids coding for Mirk (Wt), kinase-inactive Mirk (YF) or MITR. Data courtesy of laboratory of Dr. Eileen Friedman, Upstate Medical University, New York.



Immunofluorescence staining of MITR (HDAC9) for a compartmentalization study in undifferentiated transfected C2C12 myoblasts. Data courtesy of laboratory of Dr. Eileen Friedman. Dept of Pathology, Upstate Medical University, State University of New York.



IHC analysis of FFPE human hepatocarcinoma stained with HDAC9 antibody

## Description

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene has sequence homology to members of the histone deacetylase family. This gene is orthologous to the *Xenopus* and mouse MITR genes. The MITR protein lacks the histone deacetylase catalytic domain. It represses MEF2 activity through recruitment of multicomponent corepressor complexes that include CtBP and HDACs. This encoded protein may play a role in hematopoiesis. Multiple alternatively spliced transcripts have been described for this gene but the full-length nature of some of them has not been determined.

## Application Notes

Titration of the HDAC9 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

A portion of amino acids 2-32 from the human protein was used as the immunogen for this HDAC9 antibody.

## Storage

Aliquot the HDAC9 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

