


hnRNP A1 Antibody / HNRNPA1 (R32607)


Catalog No.	Formulation	Size
R32607	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P09651
Localization	Nuclear, cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml Immunofluorescence (FFPE) : 5ug/ml Flow Cytometry : 1-3ug/million cells
Limitations	This hnRNP A1 antibody is available for research use only.




Immunofluorescent staining of FFPE human rectal cancer with hnRNP A1 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH8 EDTA for 20 min.




IHC testing of human intestinal cancer with hnRNP A1 antibody at 1ug/ml. HIER: steam section in pH6 citrate buffer for 20 min.




IHC testing of human placental tissue with hnRNP A1 antibody at 1ug/ml. HIER: steam section in pH6 citrate buffer for 20 min.




IHC testing of mouse intestine tissue with hnRNP A1 antibody at 1ug/ml. HIER: steam section in pH6 citrate buffer for 20 min.




IHC testing of mouse kidney tissue with hnRNP A1 antibody at 1ug/ml. HIER: steam section in pH6 citrate buffer for 20 min.




IHC testing of rat kidney tissue with hnRNP A1 antibody at 1ug/ml. HIER: steam section in pH6 citrate buffer for 20 min.




Western blot testing of human 1) HepG2, 2) Daudi, 3) MOLT4 and 4) HL60 cell lysate with hnRNP A1 antibody. Expected molecular weight: 29-39 kDa (multiple isoforms).



Immunofluorescent staining of FFPE human A431 cells with hnRNP A1 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Immunofluorescent staining of FFPE human U-2 OS cells with hnRNP A1 antibody (red) and DAPI nuclear stain (blue). HIER: steam section in pH6 citrate buffer for 20 min.



Flow cytometry testing of human K562 cells with hnRNP A1 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= hnRNP A1 antibody.

Description

Heterogeneous nuclear ribonucleoprotein A1 is a protein that in humans is encoded by the HNRNPA1 gene. This gene encodes a member of a family of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs), which are RNA-binding proteins that associate with pre-mRNAs in the nucleus and influence pre-mRNA processing, as well as other aspects of mRNA metabolism and transport. The protein encoded by this gene is one of the most abundant core proteins of hnRNP complexes and plays a key role in the regulation of alternative splicing. Mutations in this gene have been observed in individuals with amyotrophic lateral sclerosis 20. Multiple alternatively spliced transcript variants have been found. There are numerous pseudogenes of this gene distributed throughout the genome.

Application Notes

Optimal dilution of the hnRNP A1 antibody should be determined by the researcher.

Immunogen

Amino acids 8-42 (KEPEQLRKLFIGGLSFETTDSELSRSHFEQWGT LTD) were used as the immunogen for the hnRNP A1 antibody.

Storage

After reconstitution, the hnRNP A1 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.