


## HSP60 Antibody / HSPD1 (N1068)


| Catalog No. | Formulation                                                             | Size   |
|-------------|-------------------------------------------------------------------------|--------|
| N1068-100UG | 0.5 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| N1068-25UG  | 0.5 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 25 ug  |

[Bulk quote request](#)


|                    |                                                                   |
|--------------------|-------------------------------------------------------------------|
| Species Reactivity | Human                                                             |
| Format             | Purified                                                          |
| Clonality          | Polyclonal (rabbit origin)                                        |
| Isotype            | Rabbit IgG                                                        |
| Purity             | Protein A affinity chromatography                                 |
| Buffer             | 1X PBS, pH 7.4                                                    |
| Gene ID            | 3329                                                              |
| Applications       | Western Blot : 1-3ug/ml<br>Immunohistochemistry (FFPE) : 5-7ug/ml |
| Limitations        | This <b>HSP60 antibody</b> is available for research use only.    |



Western blot testing of human samples with HSP60 antibody at 1ug/ml.



Western blot testing of mouse samples with HSP60 antibody at 1ug/ml.



IHC staining of FFPE human small intestine with HSP60 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 20 min and allow to cool before testing.

## Description

First described in relation to heat shock, HSPs are now known to be produced in response to additional stressful conditions including cold and UV light, as well as during wound healing. Heat shock protein 60 (HSP60) is responsible for protein transportation from the cytoplasm to the mitochondrial matrix and the subsequent refolding of the transported proteins. HSP60 also functions as a molecular chaperone, or chaperonin, assisting linear amino acid chains to fold into their three-dimensional structure.

## Application Notes

Provided assay concentrations are suggestions only, HSP60 antibody titration may be required for optimal results.

## Immunogen

A recombinant protein fragment from human HSP60 was used as the immunogen for this antibody.

## Storage

Aliquot the HSP60 antibody and store frozen at -20oC or colder to avoid repeated freeze-thaw cycles.

## Alternate Names

Chaperonin 60-KD (CPN60), GROEL, HLD4, HSP65, HSPD1, HuCHA60, Mitochondrial matrix protein P1, Hsp60s1, Spastic paraplegia 13 (SPG13), HSP60 antibody