


HTT Antibody / Huntingtin (RQ8713)


Catalog No.	Formulation	Size
RQ8713	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity chromatography
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	P42858
Localization	Cytoplasm, Nucleus
Applications	Immunohistochemistry (FFPE) : 2-5ug/ml ELISA : 0.1-0.5ug/ml
Limitations	This HTT antibody is available for research use only.




IHC staining of FFPE human lung cancer tissue with HTT antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human liver cancer tissue with HTT antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human endometrioid adenocarcinoma tissue with HTT antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human colon adenocarcinoma tissue with HTT antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human breast cancer tissue with HTT antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human cervix squamous cell carcinoma tissue with HTT antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human placental tissue with HTT antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

The huntingtin gene, also called HTT or HD (Huntington disease) gene, is the IT15 (interesting transcript 15) gene which codes for a protein called the huntingtin protein. It is mapped to 4p16.3. The protein has no sequence homology with other proteins and is highly expressed in neurons and tests in humans and rodents. HTT upregulates the expression of Brain Derived Neurotrophic Factor (BDNF) at the transcription level, and this gene is required for normal development. The HTT protein is involved in vesicle trafficking as it interacts with HIP1, a clathrin-binding protein, to mediate endocytosis, the absorption of materials into a cell. HTT was also visualized as punctate staining likely to represent nerve endings. What's more, wildtype HTT may function in the nucleus in the assembly of nuclear matrix-bound protein complexes involved with transcriptional repression and RNA processing.

Application Notes

Optimal dilution of the HTT antibody should be determined by the researcher.

Immunogen

An E.coli-derived human recombinant protein (amino acids R3079-C3142) was used as the immunogen for the HTT antibody.

Storage

After reconstitution, the HTT antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.