


ICAM3 Antibody / Intercellular adhesion molecule 3 / CD50 [clone 28I48] (RQ8387)

| Catalog No. | Formulation | Size |
|-------------|----------------------------------------------------------------------------|--------|
| RQ8387 | Antibody in PBS with 0.02% sodium azide, 50% glycerol and 0.4-0.5mg/ml BSA | 100 ul |

Recombinant RABBIT MONOCLONAL

Bulk quote request

| Availability | 1-3 business days |
|--------------------|---------------------------------------------------------|
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Recombinant Rabbit Monoclonal |
| Isotype | Rabbit IgG |
| Clone Name | 28148 |
| Purity | Affinity chromatography |
| UniProt | P32942 |
| Applications | Western Blot : 1:500-1:2000 |
| Limitations | This ICAM3 antibody is available for research use only. |



Western blot testing of human Ramos cell lysate with ICAM3 antibody. Observed molecular weight: 60~120 kDa depending on glycosylation level.

Description

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM3 is a ligand for integrin alpha-D/beta-2. In association with integrin alpha-L/beta-2, contributes to apoptotic neutrophil phagocytosis by macrophages. [UniProt]

Application Notes

Optimal dilution of the ICAM3 antibody should be determined by the researcher.

Immunogen

A synthetic peptide derived from human ICAM3 / CD50 was used as the immunogen for the ICAM3 antibody.

Storage

Store the ICAM3 antibody at -20oC.