


IgA Antibody [clone IA761] (V2619CF488)


Catalog No.	Formulation	Size
V2619CF488-100T	500 ul at 0.1 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 Tests

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	CF488 Conjugate
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	IA761
Purity	Protein G affinity chromatography
UniProt	P01876
Localization	Cell surface, cytoplasmic, secreted
Applications	Flow Cytometry : 5ul/test/10 ⁶ cells in 0.1ml Immunofluorescence : 1-2ug/ml
Limitations	This IgA antibody is available for research use only.



Immunofluorescent staining of PFA-fixed human Raji cells with CF488-conjugated IgA antibody (clone IA761, green) and Reddot nuclear stain (red).



SDS-PAGE analysis of purified, BSA-free IgA antibody (clone IA761) as confirmation of integrity and purity.

Description

This MAb is specific to heavy chain of IgA and shows minimal cross-reaction with heavy chains of other immunoglobulins. It is reactive with all subclasses of Alpha heavy chain. Immunoglobulins are four-chain, Y-shaped, monomeric structures comprised of two identical heavy chains and two identical light chains held together through inter-chain disulfide bonds. The chains form two domains, the Fab (antigen binding) fragment and the Fc (constant) fragment. Immunoglobulin A (IgA) is the main protein of the mucosal immune system. It is generated by B-cells in gut-associated lymphoid tissues. Daily production of IgA exceeds that of any of the other immunoglobulins. IgA exists mainly in dimers but can also exist as polymers or as monomers. Dimers and polymers contain a joining (J) chain that can be bound by the polymeric immunoglobulin receptor (pIgR) for transportation of the molecule to mucosal surfaces. The most common feature of plasmacytomas, and certain non-Hodgkin's lymphomas is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is clonal and therefore malignant.

Application Notes

Optimal dilution of the IgA antibody should be determined by the researcher.

Immunogen

Purified human immunoglobulin alpha heavy chain was used as the immunogen for the Anti-IgA antibody.

Storage

Store the IgA antibody at 2-8°C, protected from light.