


## IL12 p40 Antibody / IL12B (RQ4076)

Catalog No.	Formulation	Size
RQ4076	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

## **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human, Mouse
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P29460
Applications	Western Blot: 0.5-1ug/ml Direct ELISA: 0.1-0.5ug/ml
Limitations	This IL12 p40 antibody is available for research use only.



Western blot testing of 1) mouse NIH3T3 and 2) human A549 cell lysate with IL12 p40 antibody at 0.5 $\mu$ ml. Predicted molecular weight ~37 kDa, observed molecular weight ~40 kDa.

# **Description**

Subunit beta of interleukin 12 (IL12B), also known as interleukin 12B, or interleukin-12 subunit p40, is a subunit of human interleukin 12. It is mapped to 5q33.3. This cytokine is expressed by activated macrophages that serve as an essential inducer of Th1 cells development. This cytokine has been found to be important for sustaining a sufficient number of memory/effector Th1 cells to mediate long-term protection to an intracellular pathogen. Overexpression of this gene was observed in the central nervous system of patients with multiple sclerosis (MS), suggesting a role of this cytokine in the pathogenesis of the disease.

## **Application Notes**

Optimal dilution of the IL12 p40 antibody should be determined by the researcher.

#### **Immunogen**

A recombinant human partial protein corresponding to amino acids I23-E253 was used as the immunogen for the IL12 p40 antibody.

### **Storage**

After reconstitution, the IL12 p40 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.